



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68, C07H 21/04	A1	(11) International Publication Number: WO 97/14811 (43) International Publication Date: 24 April 1997 (24.04.97)
<p>(21) International Application Number: PCT/US96/16085</p> <p>(22) International Filing Date: 7 October 1996 (07.10.96)</p> <p>(30) Priority Data: 08/546,130 20 October 1995 (20.10.95) US</p> <p>(71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 22nd floor, 300 Lakeside Drive, Oakland, CA 94612 (US).</p> <p>(72) Inventors: GRAY, Joe, W.; 1921 11th Avenue, San Francisco, CA 94116 (US). COLLINS, Colin; 36 Shannan Lane, San Rafael, CA 94901 (US). PINKEL, Daniel; 31 Manzanita Court, Walnut Creek, CA 94595 (US). KALLIONIEMI, Olli-Pekka; Liljankuja 4, FIN-33300 Tampere (FI). TANNER, Minna, M.; Tanhuankatu 48 C 38, FIN-33560 Tampere (FI).</p> <p>(74) Agents: HUNTER, Tom et al.; Townsend and Townsend and Crew L.L.P., 8th floor, Two Embarcadero Center, San Francisco, CA 94111 (US).</p>		<p>(81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report.</i></p>
<p>(54) Title: AMPLIFICATIONS OF CHROMOSOMAL REGION 20q13 AS A PROGNOSTIC INDICATOR IN BREAST CANCER</p> <p>(57) Abstract</p> <p>The present invention relates to <i>in situ</i> hybridization methods for the identification of new chromosomal abnormalities associated with various diseases. In particular, it provides probes which are specific to a region of amplification in chromosome 20.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

NOT AVAILABLE COPY

AMPLIFICATIONS OF CHROMOSOMAL REGION 20q13 AS A PROGNOSTIC INDICATOR IN BREAST CANCER

BACKGROUND OF THE INVENTION

This invention pertains to the field to the field of cytogenetics. More particularly this invention pertains to the identification of a amplification at about 20q13 that is a good prognositic indicator of various cancers. In addition, this invention provides a number of probes specific for the 20q13 amplicon.

This invention was made with Government support under Grant Nos. CA-528207 and CA-44768 awarded by the National Institutes of Health and Contract Nos. DE-AC-03-76SF00098, W-7405-ENG-48 and W-7405-ENG-36 awarded by the U.S. Department of Energy. The Government has certain rights in this invention.

Chromosome abnormalities are often associated with genetic disorders, degenerative diseases, and cancer. In particular, the deletion or multiplication of copies of whole chromosomes or chromosomal segments, and higher level amplifications of specific regions of the genome are common occurrences in cancer. See, for example Smith, *et al.*, *Breast Cancer Res. Treat.*, 18: Suppl. 1: 5-14 (1991, van de Vijer & Nusse, *Biochim. Biophys. Acta.* 1072: 33-50 (1991), Sato, *et al.*, *Cancer. Res.*, 50: 7184-7189 (1990). In fact, the amplification and deletion of DNA sequences containing proto-oncogenes and tumor-suppressor genes, respectively, are frequently characteristic of tumorigenesis. Dutrillaux, *et al.*, *Cancer Genet. Cytogenet.*, 49: 203-217 (1990). Clearly the identification of amplified and deleted regions and the cloning of the genes involved is crucial both to the study of tumorigenesis and to the development of cancer diagnostics.

The detection of amplified or deleted chromosomal regions has traditionally been done by cytogenetics. Because of the complex packing of DNA into the chromosomes, resolution of cytogenetic techniques has been limited to regions larger than about 10 Mb; approximately the width of a band in Giemsa-stained chromosomes. In complex karyotypes with multiple translocations and other genetic changes, traditional cytogenetic analysis is of little utility because karyotype information is lacking or cannot be interpreted. Teyssier, J.R., *Cancer Gener. Cytogenet.*, 37: 103 (1989). Furthermore

conventional cytogenetic banding analysis is time consuming, labor intensive, and frequently difficult or impossible.

More recently, cloned probes have been used to assess the amount of a given DNA sequence in a chromosome by Southern blotting. This method is effective even if the genome is heavily rearranged so as to eliminate useful karyotype information. However, Southern blotting only gives a rough estimate of the copy number of a DNA sequence, and does not give any information about the localization of that sequence within the chromosome.

Comparative genomic hybridization (CGH) is a more recent approach to identify the presence and localization of amplified/deleted sequences. See Kallioniemi, *et al.*, *Science*, 258: 818 (1992). CGH, like Southern blotting, reveals amplifications and deletions irrespective of genome rearrangement. Additionally, CGH provides a more quantitative estimate of copy number than Southern blotting, and moreover also provides information of the localization of the amplified or deleted sequence in the normal chromosome.

Using CGH, the chromosomal 20q13 region has been identified as a region that is frequently amplified in cancers (*see, e.g.* copending application U.S.S.N. 08/132,808, filed on October 6, 1993). Initial analysis of this region in breast cancer cell lines identified a region approximately 2 Mb on chromosome 20 that is consistently amplified.

SUMMARY OF THE INVENTION

The present invention relates to the identification of a narrow region (about 600 kb) within a 2 Mb amplicon located at about chromosome 20q13 (more precisely at 20q13.2) that is consistently amplified in primary tumors. In addition this invention provides a contig (a series of clones that contiguously spans this amplicon) as well as sequence information for a large number of exons and cDNAs located within the contig. The contig or components thereof can be used to prepare probes specific for the amplicon. The probes can be used to detect chromosomal abnormalities at 20q13.

Thus, in one embodiment, this invention provides a method of detecting a chromosome abnormality (*e.g.*, an amplification or a deletion) at about position FLpter 0.825 on human chromosome 20 (20q13.2). The method involves contacting a chromosome sample from a patient with a composition consisting essentially of one or

BEST AVAILABLE COPY

more labeled nucleic acid probes each of which binds selectively to a target polynucleotide sequence at about position FLpter 0.825 on human chromosome 20 under conditions in which the probe forms a stable hybridization complex with the target sequence; and detecting the hybridization complex. The step of detecting the hybridization complex can involve determining the copy number of the target sequence. The probe preferably comprises a nucleic acid that specifically hybridizes under stringent conditions to a nucleic acid selected from the nucleic acids listed in Table 1 or Table 2. Even more preferably, the probe is one or more nucleic acids selected from the nucleic acids listed in Table 1 or Table 2. The probe is preferably labeled, and is more preferably labeled with digoxigenin or biotin. In one embodiment, the hybridization complex is detected in interphase nuclei in the sample. Detection is preferably carried out by detecting a fluorescent label (e.g., FITC, fluorescein, or Texas Red). The method can further involve contacting the sample with a reference probe which binds selectively to a chromosome 20 centromere.

In another embodiment, this invention provides for probes that specifically bind to the 20q13 amplicon. Thus, this invention provides for a composition comprising a labeled nucleic acid probe which binds selectively to a target polynucleotide sequence at about FLpter 0.825 on human chromosome 20. The probes comprise one or more nucleic acids selected from the group consisting of the nucleic acids listed in Table 1 or Table 2. In a preferred embodiment, the probes are labelled with digoxigenin or biotin.

This invention also provides for kits for the detection of a chromosomal abnormality at about position FLpter 0.825 on human chromosome 20. The kits include a compartment which contains a labeled nucleic acid probe which binds selectively to a target polynucleotide sequence at about FLpter 0.825 on human chromosome 20. The probe preferably includes at least one nucleic acid that specifically hybridizes under stringent conditions to a nucleic acid selected the nucleic acids listed in Table 1 or Table 2. Even more preferably, the probes comprise one or more nucleic acids selected from the nucleic acids listed in Table 1 or Table 2. In a preferred embodiment, the probes are labelled with digoxigenin or biotin. The kit may further include a reference probe specific to a sequence in the centromere of chromosome 20.

BEST AVAILABLE COPY

Definitions

A "chromosome sample" as used herein refers to a tissue or cell sample prepared for standard *in situ* hybridization methods described below. The sample is prepared such that individual chromosomes remain substantially intact and typically comprises metaphase spreads or interphase nuclei prepared according to standard techniques.

"Nucleic acid" refers to a deoxyribonucleotide or ribonucleotide polymer in either single- or double-stranded form, and unless otherwise limited, would encompass known analogs of natural nucleotides that can function in a similar manner as naturally occurring nucleotides.

"Subsequence" refers to a sequence of nucleic acids that comprise a part of a longer sequence of nucleic acids.

A "probe" or a "nucleic acid probe", as used herein, is defined to be a collection of one or more nucleic acid fragments whose hybridization to a target can be detected. The probe is labeled as described below so that its binding to the target can be detected. The probe is produced from a source of nucleic acids from one or more particular (preselected) portions of the genome, for example one or more clones, an isolated whole chromosome or chromosome fragment, or a collection of polymerase chain reaction (PCR) amplification products. The probes of the present invention are produced from nucleic acids found in the 20q13 amplicon as described herein. The probe may be processed in some manner, for example, by blocking or removal of repetitive nucleic acids or enrichment with unique nucleic acids. Thus the word "probe" may be used herein to refer not only to the detectable nucleic acids, but to the detectable nucleic acids in the form in which they are applied to the target, for example, with the blocking nucleic acids, *etc.* The blocking nucleic acid may also be referred to separately. What "probe" refers to specifically is clear from the context in which the word is used.

"Hybridizing" refers the binding of two single stranded nucleic acids via complementary base pairing.

"Bind(s) substantially" or "binds specifically" or "binds selectively" or "hybridizing specifically to" refers to complementary hybridization between an oligonucleotide and a target sequence and embraces minor mismatches that can be accommodated by reducing the stringency of the hybridization media to achieve the

NOT AVAILABLE COPY

desired detection of the target polynucleotide sequence. These terms also refer to the binding, duplexing, or hybridizing of a molecule only to a particular nucleotide sequence under stringent conditions when that sequence is present in a complex mixture (e.g., total cellular) DNA or RNA. The term "stringent conditions" refers to conditions under which a probe will hybridize to its target subsequence, but to no other sequences.

Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures.

Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength, pH, and nucleic acid concentration) at which 50% of the probes complementary to the target sequence hybridize to the target sequence at equilibrium. Typically, stringent conditions will be those in which the salt concentration is at least about 0.02 Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 60°C for short probes. Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide.

One of skill will recognize that the precise sequence of the particular probes described herein can be modified to a certain degree to produce probes that are "substantially identical" to the disclosed probes, but retain the ability to bind substantially to the target sequences. Such modifications are specifically covered by reference to the individual probes herein. The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 90% sequence identity, more preferably at least 95%, compared to a reference sequence using the methods described below using standard parameters.

Two nucleic acid sequences are said to be "identical" if the sequence of nucleotides in the two sequences is the same when aligned for maximum correspondence as described below. The term "complementary to" is used herein to mean that the complementary sequence is identical to all or a portion of a reference polynucleotide sequence.

Sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two sequences over a "comparison window" to identify and compare local regions of sequence similarity. A "comparison window", as used herein, refers to a segment of at least about 20 contiguous positions, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be

compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman *Adv. Appl. Math.* 2: 482 (1981), by the homology alignment algorithm of Needleman and Wunsch *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson and Lipman *Proc. Natl. Acad. Sci. (U.S.A.)* 85: 2444 (1988), by computerized implementations of these algorithms.

"Percentage of sequence identity" is determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

Another indication that nucleotide sequences are substantially identical is if two molecules hybridize to the same sequence under stringent conditions. Stringent conditions are sequence dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically, stringent conditions will be those in which the salt concentration is about 0.02 molar or lower at pH 7 and the temperature is at least about 60°C.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1(A) shows disease-free survival of 129 breast cancer patients

BEST AVAILABLE COPY

according to the level of 20q13 amplification. Patients with tumors having high level 20q13 amplification have a shorter disease-free survival ($p=0.04$ by Mantel-Cox test) compared to those having no or low level amplification.

Figure 1(B) Shows the same disease-free survival difference of Figure 4(A) in the sub-group of 79 axillary node-negative patients ($p=0.0022$ by Mantel-Cox test).

Figure 2 shows a comparison of 20q13 amplification detected by FISH in a primary breast carcinoma and its metastasis from a 29-year patient. A low level amplification of 20q13 (20q13 compared to 20p reference probe) was found in the primary tumor. The metastasis, which appeared 8 months after mastectomy, shows a high level amplification of the chromosome 20q13 region. The overall copy number of chromosome 20 (20p reference probe) remained unchanged. Each data point represents gene copy numbers in individual tumor cells analyzed.

Figure 3 shows a graphical representation of the molecular cytogenetic mapping and subsequent cloning of the 20q13.2 amplicon. Genetic distance is indicated in centiMorgans (cM). The thick black bar represents the region of highest level amplification in the breast cancer cell line BT474 and covers a region of about 1.5 Mb. P1 and BAC clones are represented as short horizontal lines and YAC clones as heavier horizontal lines. Not all YAC and P1 clones are shown. YACs 957f3, 782c9, 931h12, and 901b12 are truncated. Sequence tagged sites (STSs) appear as thin vertical lines and open circles indicate that a given YAC has been tested for and is positive for a given STS. Not all STSs have been tested on all YACs. The interval from which more than 100 exons have been trapped is represented as a filled box. The 600 kb interval spanning the region of highest amplification level in primary tumors is represented by the filled black box (labeled Sequence). The lower part of the figure shows the levels of amplification in two primary tumors that further narrow the region of highest amplification to within about 600 kb.

Figure 4 provides a higher resolution map of the amplicon core as defined in primary tumors.

DETAILED DESCRIPTION

This invention provides a number of nucleic acid probes useful for the detection of chromosomal abnormalities at 20q13. Studies using comparative genomic

BEST AVAILABLE COPY

hybridization (CGH) have shown that a region at chromosome 20q13 is increased in copy number frequently in cancers of the breast (~30%), ovary (~15%), bladder (~30%), head and neck (~75%) and colon (~80%). This suggests the presence of one or more genes that contribute to the progression of several solid tumors are located at 20q13.

Gene amplification is one mechanism by which dominantly acting oncogenes are overexpressed, allowing tumors to acquire novel growth characteristics and/or resistance to chemotherapeutic agents. Loci implicated in human breast cancer progression and amplified in 10-25% of primary breast carcinomas include the *erbB-2* locus (Lupu *et al.*, *Breast Cancer Res. Treat.*, 27: 83 (1993), Slamon *et al.* *Science*, 235: 177-182 (1987), Heiskanen *et al.* *Biotechniques*, 17: 928 (1994)) at 17q12, cyclin-D (Mahadevan *et al.*, *Science*, 255: 1253-1255 (1993), Gillett *et al.*, *Canc. Res.*, 54: 1812 (1994)) at 11q13 and MYC (Gaffey *et al.*, *Mod. Pathol.*, 6: 654 (1993)) at 8q34.

Pangenomic surveys using comparative genomic hybridization (CGH) recently identified about 20 novel regions of increased copy number in breast cancer (Kallioniemi *et al.*, *Genomics*, 20: 125-128 (1994)). One of these loci, band 20q13, was amplified in 18% of primary tumors and 40% of cell lines (Kallioniemi *et al.*, *Genomics*, 20: 125-128 (1994)). More recently, this same region was found amplified in 15% of ovarian, 80% of bladder and 80% of colorectal tumors. The resolution of CGH is limited to 5-10 Mb. Thus, FISH was performed using locus specific probes to confirm the CGH data and precisely map the region of amplification.

The 20q13 region has been analyzed in breast cancer at the molecular level and a region, approximately 600 kb wide, that is consistently amplified was, identified, as described herein. Moreover, as shown herein, the importance of this amplification in breast cancer is indicated by the strong association between amplification and decreased patient survival and increased tumor proliferation (specifically, increased fraction of cells in S-phase).

In particular, as explained in detail in Example 1, high-level 20q13 amplification was associated ($p=0.0022$) with poor disease free survival in node-negative patients, compared to cases with no or low-level amplification (Figure 1). Survival of patients with moderately amplified tumors did not differ significantly from those without amplification. Without being bound to a particular theory, it is suggested that an explanation for this observation may be that low level amplification precedes high level amplification. In this regard, it may be significant that one patient developed a local

BEST AVAILABLE COPY

metastasis with high-level 20q13.2 amplification 8 month after resection of a primary tumor with low level amplification (Figure 3).

The 20q13 amplification was associated with high histologic grade of the tumors. This correlation was seen in both moderately and highly amplified tumors.

There was also a correlation ($p=0.0085$) between high level amplification of a region complementary to a particular probe, RMC20C001 (Tanner *et al.*, *Cancer Res.*, 54: 4257-4260 (1994)), and cell proliferation, measured by the fraction of cells in S-phase (Figure 4). This finding is important because it identifies a phenotype that can be scored in functional assays, without knowing the mechanism underlying the increased S-phase fraction. The 20q13 amplification did not correlate with the age of the patient, primary tumor size, axillary nodal or steroid hormone-receptor status.

This work localized the 20q13.2 amplicon to an interval of approximately 2 Mb. Furthermore, it suggests that high-level amplification, found in 7% of the tumors, confers an aggressive phenotype on the tumor, adversely affecting clinical outcome. Low level amplification (22% of primary tumors) was associated with pathological features typical of aggressive tumors (high histologic grade, aneuploidy and cell proliferation) but not patient prognosis.

In addition, it is shown herein that the 20q13 amplicon (more precisely the 20q13.2 amplicon) is one of three separate co-amplified loci on human chromosome 20 that are packaged together throughout the genomes of some primary tumors and breast cancer cell lines. No known oncogenes map in the 20q13.2 amplicon.

Identification of 20q13 Amplicon Probes.

Initially, a paucity of available molecular cytogenetic probes dictated that FISH probes be generated by the random selection of cosmids from a chromosome 20 specific library, LA20NC01, and map them to chromosome 20 by digital imaging microscopy. Approximately 46 cosmids, spanning the 70 Mb chromosome, were isolated for which fractional length measurements (FLpter) and band assignments were obtained. Twenty six of the cosmids were used to assay copy number in the breast cancer cell line BT474 by interphase FISH analysis. Copy number was determined by counting hybridization signals in interphase nuclei. This analysis revealed that cosmid RMC20C001 (Flpter, 0.824; 20q13.2), described by Stokke *et al.*, *Genomics*, 26: 134-

BEST AVAILABLE COPY

137 (1995), defined the highest-level amplification (~60 copies/cell) in BT474 cells (Tanner *et al.*, *Cancer Res.*, 54: 4257-4260 (1994)).

P1 clones containing genetically mapped sequences were selected from 20q13.2 and used as FISH probes to confirm and further define the region of amplification. Other P1 clones were selected for candidate oncogenes broadly localized to the 20q13.2 region (Flpter, 0.81-0.84). These were selected from the DuPont P1 library (Shepherd, *et al.*, *Proc. Natl. Acad. Sci. USA*, 92: 2629 (1994), available commercially from Genome Systems), by PCR (Saiki *et al.*, *Science*, 230: 1350 (1985)) using primer pairs developed in the 3' untranslated region of each candidate gene. Gene specific P1 clones were obtained for, protein tyrosine phosphatase (PTPN1, Flpter 0.78), melanocortin 3 receptor (MC3R, Flpter 0.81), phosphoenolpyruvate carboxy kinase (PCK1, Flpter 0.85), zinc finger protein 8 (ZNF8, Flpter 0.93), guanine nucleotide-binding protein (GNAS 1, Flpter .873), src-oncogene (SRC, Flpter 0.669), topoisomerase 1 (TOP1, Flpter 0.675), the bcl-2 related gene bcl-x (Flpter 0.526) and the transcription factor E2F-1 (FLpter 0.541). Each clone was mapped by digital imaging microscopy and assigned Flpter values. Five of these genes (SRC, TOPO1, GNAS1, E2F-1 and BC1-x) were excluded as candidate oncogenes in the amplicon because they mapped well outside the critical region at Flpter 0.81-0.84. Three genes (PTPNR1, PCK-1 and MC3R) localized close enough to the critical region to warrant further investigation.

Interphase FISH on 14 breast cancer cell lines and 36 primary tumors using 24 cosmid and 3 gene specific P1 (PTPNRL, PCK-1 and MC3R) probes found high level amplification in 35% (5/14) of breast cancer cell lines and 8% (3/36) of primary tumors with one or more probe. The region with the highest copy number in 4/5 of the cell lines and 3/3 primary tumors was defined by the cosmid RMC20C001. This indicated that PTPNR1, PCK1 and MC3R could also be excluded as candidates for oncogenes in the amplicon and, moreover, narrowed the critical region from 10 Mb to 1.5-2.0 Mb (*see*, Tanner *et al.*, *Cancer Res.*, 54: 4257-4260 (1994)).

Because probe RMC20C001 detected high-level (3 to 10-fold) 20q13.2 amplification in 35% of cell lines and 8% of primary tumors it was used to (1) define the prevalence of amplification in an expanded tumor population, (2) assess the frequency and level of amplification in these tumors, (3) evaluate the association of the 20q13.2 amplicon with pathological and biological features, (4) determine if a relationship exists

BEST AVAILABLE COPY

between 20q13 amplification and clinical outcome and (5) assess 20q13 amplification in metastatic breast tumors.

As detailed in Example 1, fluorescent *in situ* hybridization (FISH) with RMC20C001 was used to assess 20q13.2 amplification in 132 primary and 11 recurrent breast tumors. The absolute copy number (mean number of hybridization signals per cell) and the level of amplification (mean number of signals relative to the p-arm reference probe) were determined. Two types of amplification were found: Low level amplification (1.5-3 fold with FISH signals dispersed throughout the tumor nuclei) and high level amplification (>3 fold with tightly clustered FISH signals). Low level 20q13.2 amplification was found in 29 of the 132 primary tumors (22%), whereas nine cases (6.8%) showed high level amplification.

RMC20C001 and four flanking P1 probes (MC3R, PCK, RMC20C026, and RMC20C030) were used to study the extent of DNA amplification in highly amplified tumors. Only RMC20C001 was consistently amplified in all tumors. This finding confirmed that the region of common amplification is within a 2 Mb interval flanked by but not including PCK-1 and MC3R.

A physical map was assembled to further localize the minimum common region of amplification and to isolate the postulated oncogene(s). The DuPont P1 library (Shepherd *et al. Proc. Natl. Acad. Sci. USA*, 91: 2629 (1994)) was screened for STSs likely to map in band 20q13.2. P1 clones at the loci D20S102, D20S100, D20S120, D20S183, D20S480, D20S211 were isolated, and FISH localized each to 20q13.2. Interphase FISH analysis was then performed in the breast cancer cell line BT474 to assess the amplification level at each locus. The loci D20S100-D20S120-D20S183-D20S480-D20S211 were highly amplified in the BT474 cell line, whereas D20S102 detected only low level amplification. Therefore, 5 STSs, spanning 5 cM, were localized within the 20q13.2 amplicon and were utilized to screen the CEPH megaYAC library.

CEPH megaYAC library screening and computer searches of public databases revealed D20S120-D20S183-D20S480-D20S211 to be linked on each of three megaYAC clones y820f5, 773h10, and 931h6 (Figure 3). Moreover, screening the CEPH megaYAC library with STSs generated from the ends of cosmids RMC20C001, RMC20C30 and RMC20C028 localized RMC20C001 to each of the same three YAC clones. It was estimated, based on the size of the smallest of these YAC clones, that D20S120-D20S 183-RMC20C001-D20S480-D20S211 map into an interval of less than

BEST AVAILABLE COPY

1.1 Mb. D20S100 was localized 300 kb distal to D20S120 by interphase FISH and to YAC901b12 by STS mapping. The combined STS data made it possible to construct a 12 member YAC contig which spans roughly 4 Mb encompassing the 1.5 Mb amplicon and containing the loci RMC20C030-PCK1-RMC20C001-MC3R-RMC20C026. Each
5 YAC was mapped by FISH to confirm localization to 20q13.2 and to check for chimerism. Five clonal isolates of each YAC were sized by pulsed field gel electrophoresis (PFGE). None of the YACs are chimeric, however, several are highly unstable.

The YAC contig served as a framework from which to construct a 2 MbP1
10 contig spanning the 20Q13 amplicon. P1 clones offered numerous advantages over YAC clones including (1) stability, (2) a chimeric frequency of less than 1%, (3) DNA isolation by standard miniprep procedures, (4) they make ideal FISH probes, (5) the ends can be sequenced directly, (6) engineered $\gamma\delta$ transposons carrying bidirectional primer binding sites can be integrated at any position in the cloned DNA (Strathmann *et al.*,
15 *Proc. Natl. Acad. Sci. USA*, 88: 1247 (1991)) (7) P1 clones are the templates for sequencing the human and Drosophila genomes at the LBNL HGC (Palazzolo *et al.* DOE Human Genome Program, Contractor-Grantee Workshop IV. Santa Fe, New Mexico, November 13-17 1994).

About 90 P1 clones were isolated by screening the DuPont P1 library
20 either by PCR or filter hybridization. For PCR based screening, more than 22 novel STSs were created by two methods. In the first method, the ends of P1 clones localized to the amplicon were sequenced, STSs developed, and the P1 library screened for walking clones. In the second approach inter-Alu PCR (Nelson *et al.*, 86: 6686-6690 (1989)) was performed on YACs spanning the amplicon and the products cloned and
25 sequenced for STS creation. In the filter based hybridization scheme P1 clones were obtained by performing inter-Alu PCR on YACs spanning the amplicon, radio-labeling the products and hybridizing these against filters containing a gridded array of the P1 library. Finally, to close gaps a human genomic bacterial artificial chromosome (BAC) library (Shizuya *et al.* *Proc. Natl. Acad. Sci. USA*, 89: 8794 (1992), commercially
30 available from Research Genetics, Huntsville, Alabama, USA) was screened by PCR. These methods combined to produce more than 100 P1 and BAC clones were localized to 20q13.2 by FISH. STS content mapping, fingerprinting, and free-chromatin fish

BEST AVAILABLE COPY

(Heiskanen *et al.*, *BioTechniques*, 17: 928 (1994)) were used to construct the 2 Mb contig shown in Figure 3.

Fine Mapping the 20q13.2 Amplicon in BT474

5 Clones from the 2 Mb P1 contig were used with FISH to map the level of amplification at 20q13.2 in the breast cancer cell line BT474. 35 P1 probes distributed at regular intervals along the contig were used. The resulting data indicated that the region of *highest copy number increase* in BT474 occurs between D20S100 and D20S211, an interval of approximately 1.5 Mb. P1 FISH probes, in this interval, detect
10 an average of 50 signals per interphase nuclei in BT474, while no, or only low level amplification, was detected with the P1 clones outside this region. Thus, both the proximal and distal boundaries of the amplicon were cloned.

Fine Mapping the 20q13.2 Amplicon in Primary Tumors.

15 Fine mapping the amplicon in primary tumors revealed the *minimum common region of high amplification* that is of pathobiological significance. This process is analogous to screening for informative meiosis in the narrowing of genetic intervals encoding heritable disease genes. Analysis of 132 primary tumors revealed thirty-eight primary tumors that are amplified at the RMC20C001 locus. Nine of these tumors have
20 high level amplification at the RMC20C001 locus and were further analyzed by interphase FISH with 8 P1s that span the ≈ 2 Mb contig. The minimum common region of amplification was mapped to a ≈ 600 kb interval flanked by P1 clones #3 and #12 with the highest level of amplification detected by P1 clone #38 corresponding to RMC20C001 (Figure 4).

25 The P1 clones spanning the 600 kb interval of the 20q13 amplicon are listed in Table 1 which provides a cross-reference to the DuPont P1 library described by Shepherd, *et al.*, *Proc. Natl. Acad. Sci. USA*, 92: 2629 (1994). These P1 probes are available commercially from Genetic Systems.

BEST AVAILABLE COPY

Table 1. Cross reference to probes of the DuPont P1 library (Shepherd, *et al.*, *Proc. Natl. Acad. Sci. USA*, 92: 2629 (1994) which is commercially available from Genomic Systems, St. Louis Missouri, USA). PCR primers are illustrated for amplification of Sequence tag sites for each clone. In addition PCR conditions (Mg concentration and annealing temperature) as well as PCR product size is provided. Size: PCR product Size; mM MgCl: Mg concentration; Ann: Annealing temperature; P1: P1 probe ID number; PC: DuPont Library Plate Coordinates; SCA: DuPont Library Single Clone address; Seq-forward and Seq-backward: forward and backward PCR primers, respectively.

NAME	Probe	SIZE(bp)	mM MgCl	Ann.	P1	PC	SCA	SEQ-forward	SEQ-backward
352.32		136	1.5	52	20	103-e5	1238e	TTGGCATTGGTATCAGGTAGCTG	TTGGACGACAGACGGGATTGTGTG
388.13	F1/B1	201	1.5	52	17	59g6	821g	ATCCCTCAAACTCTGCTGTAG	TGGAGCCTGGAAGCTTCTGCATC
					19	98f4	1167f		TGCACATAAAACAGCCAGC
D20S183		270	3	48	30	124g6	..	OOGGGATACCGACATIG	
					40	24h1	276h		AGCTTTACCCAAATGTTGTC
D20S211	1/2	135	1.5	52	29	119f4	1416f	TTGGAATCAATGGAGGAAAA	AGGCATAATAAACATAAAGCTGG
D20S480		300	3	55	68	100d12	1199d2	GTGGTCAACACCAATAATGG	
					41	66c1	1020c		
					42	103d9	1232d		
					67	91b2	1081b9	CAAGAICTGACCCGTCATC	GACTTCTCAGGAAGAGATCAGTG
9X SP6 hmf/hm0		165	1.5	55	7	31d11	370d		
					9	3519	418f		
11S-17	F2/B4	148	3	58	11	41b1	480b	GGCATGTACCCAGCTGAAAAATC	TCAGACACCCGGTGCAGAAATTAAG
12T-17	F2/B1	153	3	58	12	42c2	493c	CCTAAAGCTTGGTCTTAAATCTA	GTGTACAGCCGACATGTGG
28T	F1/B1	219	1.5	52	74	88d2	..	TTTGTGATGTGAGGCATC	CTTDCATCTCATTTCTATAGG
					2	12c6	137c		
					25	118c11	1413c		
					28	118c11	1413c	GCCTGTTAAGTGTCACTAGCG	CACCTCTGGTAAATGACCTTGTG
28S	F1/B3	214	1.5	55	28	118g11	1413g		
					25	118c11	1413c		
					27	118g11	1413g		
					10	36110-	4291	CGTACACCATTCGCACTTGG	GGCAGATGTATGTTGCTACGGAAC
69S		100	3	55	69	412b5	..		
					5	23c1	264c	TTCTAAGCTGTCCAGTCTTG	CTGCTGTGTTGCGAGATGG
HSCOFH032	F/B	129	1.5	55	3	12-e11	142e		
					18	77d10	921d	TTGCTGCTCTTCCGTCATGCTAG	AATGCCCTCCACTACAGGAATG
60A1		191	1.5	58	38	112d8	1139g		
					38	34d6	401d	CCCTTCAGTGTGCTTCTATTGA	GGGAGGAGGTTGTATGCAAC
820F-5A1T	F1/B1	175	1.5	48	15	53c7	830c		
					18	58b9	692b		

BEST AVAILABLE COPY

[illegible]

The BAC clones of the contig spanning the 20q13 amplicon are listed in Table 2 which provides a cross-reference to the DuPont P1 library described by Shepherd, *et al.*, *Proc. Natl. Acad. Sci. USA*, 92: 2629 (1994). These probes are available commercially from Research Genetics.

In addition, Tables 1 and 2 provide PCR primers for amplifying sequence tagged sites (STSs) for each of the P1 and BAC probes. One of skill in the art would appreciate that using plate coordinates provided above and/or the primers

Exon Trapping

Exon trapping (*see, e.g.*, Duyk *et al.*, *Proc. Natl. Acad. Sci. USA*, 87: 8995-8999 (1990) and Church *et al.*, *Nature Genetics*, 6: 98-105 (1994)) was performed on the P1 and BAC clones spanning the \approx 600 kb minimum common region of amplification and has isolated more than 200 exons.

Analysis of the exons DNA sequence revealed a number of sequence similarities (85% to 96%) to partial CDNA sequences in the expressed sequence data base (dbest) and to a *S. cerevisiae* chromosome XIV open reading frame. Each P1 clone subjected to exon trapping has produced multiple exons consistent with at least a medium density of genes.

Detection of 20q13 Abnormalities.

One of skill in the art will appreciate that the clones and sequence information provided herein can be used to detect amplifications, or other chromosomal abnormalities, at 20q13 in a biological sample. Generally the methods involve hybridization of probes that specifically bind one or more nucleic acid sequences of the target amplicon with nucleic acids present in a biological sample or derived from a biological sample.

As used herein, a biological sample is a sample of biological tissue or fluid contains cells in it is desired to screen for chromosomal abnormalities (*e.g.* amplifications of deletions). In a preferred embodiment, the biological sample is a cell or tissue suspected of being cancerous (transformed). Methods of isolating biological samples are well known to those of skill in the art and include, but are not limited to, aspirations, tissue sections, needle biopsies, and the like. Frequently the sample will be

BEST AVAILABLE COPY

a "clinical sample" which is a sample derived from a patient. It will be recognized that the term "sample" also includes supernatant (containing cells) or the cells themselves from cell cultures, cells from tissue culture and other media in which it may be desirable to detect chromosomal abnormalities.

In a preferred embodiment, a biological sample is prepared by depositing cells, either as single cell suspensions or as tissue preparation, on solid supports such as glass slides and fixed by choosing a fixative which provides the best spatial resolution of the cells and the optimal hybridization efficiency.

Selecting Probes

Any of the P1 probes listed in Table 1 or the BAC probes listed in Table 2 are suitable for use in detecting the 20q13 amplicon. Methods of preparing probes are well known to those of skill in the art (*see, e.g. Sambrook et al., Molecular Cloning: A Laboratory Manual (2nd ed.)*, Vols. 1-3, Cold Spring Harbor Laboratory, (1989) or *Current Protocols in Molecular Biology*, F. Ausubel *et al.*, ed. Greene Publishing and Wiley-Interscience, New York (1987)).

The probes are most easily prepared by combining and labeling one or more of the constructs listed in Tables 1 and 2. Prior to use, the constructs are fragmented to provide smaller nucleic acid fragments that easily penetrate the cell and hybridize to the target nucleic acid. Fragmentation can be by any of a number of methods well known to those of skill in the art. Preferred methods include treatment with a restriction enzyme to selectively cleave the molecules, or alternatively to briefly heat the nucleic acids in the presence of Mg^{2+} . Probes are preferably fragmented to an average fragment length ranging from about 50 bp to about 2000 bp, more preferably from about 100 bp to about 1000 bp and most preferably from about 150 bp to about 500 bp.

Alternatively, probes can be produced by amplifying (*e.g. via PCR*) selected subsequences from the 20q13 amplicon disclosed herein. The sequences provided herein permit one of skill to select primers that amplify sequences from one or more exons located within the 20q13 amplicon.

Particularly preferred probes include nucleic acids from probes 38, 40, and 79, which corresponds to RMC20C001.

BEST AVAILABLE COPY

One of skill will appreciate that using the sequence information and clones provided herein, one of skill in the art can isolate the same or similar probes from other human genomic libraries using routine methods (*e.g.* Southern or Northern Blots).

Labeling Probes

Methods of labeling nucleic acids are well known to those of skill in the art. Preferred labels are those that are suitable for use in *in situ* hybridization. The nucleic acid probes may be detectably labeled prior to the hybridization reaction.

Alternatively, a detectable label which binds to the hybridization product may be used.

Such detectable labels include any material having a detectable physical or chemical property and have been well-developed in the field of immunoassays.

As used herein, a "label" is any composition detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. Useful labels in the present invention include radioactive labels (*e.g.* ^{32}P , ^{125}I , ^{14}C , ^3H , and ^{35}S), fluorescent dyes (*e.g.* fluorescein, rhodamine, Texas Red, *etc.*), electron-dense reagents (*e.g.* gold), enzymes (as commonly used in an ELISA), colorimetric labels (*e.g.* colloidal gold), magnetic labels (*e.g.* DynabeadsTM), and the like. Examples of labels which are not directly detected but are detected through the use of directly detectable label include biotin and dioxigenin as well as haptens and proteins for which labeled antisera or monoclonal antibodies are available.

The particular label used is not critical to the present invention, so long as it does not interfere with the *in situ* hybridization of the stain. However, stains directly labeled with fluorescent labels (*e.g.* fluorescein-12-dUTP, Texas Red-5-dUTP, *etc.*) are preferred for chromosome hybridization.

A direct labeled probe, as used herein, is a probe to which a detectable label is attached. Because the direct label is already attached to the probe, no subsequent steps are required to associate the probe with the detectable label. In contrast, an indirect labeled probe is one which bears a moiety to which a detectable label is subsequently bound, typically after the probe is hybridized with the target nucleic acid.

In addition the label must be detectible in as low copy number as possible thereby maximizing the sensitivity of the assay and yet be detectible above any background signal. Finally, a label must be chosen that provides a highly localized signal thereby providing a high degree of spatial resolution when physically mapping the

BEST AVAILABLE COPY

stain against the chromosome. Particularly preferred fluorescent labels include fluorescein-12-dUTP and Texas Red-5-dUTP.

The labels may be coupled to the probes in a variety of means known to those of skill in the art. In a preferred embodiment the nucleic acid probes will be labeled using nick translation or random primer extension (Rigby, *et al. J. Mol. Biol.*, 113: 237 (1977) or Sambrook, *et al., Molecular Cloning - A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1985)).

One of skill in the art will appreciate that the probes of this invention need not be absolutely specific for the targeted 20q13 region of the genome. Rather, the probes are intended to produce "staining contrast". "Contrast" is quantified by the ratio of the probe intensity of the target region of the genome to that of the other portions of the genome. For example, a DNA library produced by cloning a particular chromosome (*e.g.* chromosome 7) can be used as a stain capable of staining the entire chromosome. The library contains both sequences found only on that chromosome, and sequences shared with other chromosomes. Roughly half the chromosomal DNA falls into each class. If hybridization of the whole library were capable of saturating all of the binding sites on the target chromosome, the target chromosome would be twice as bright (contrast ratio of 2) as the other chromosomes since it would contain signal from the both the specific and the shared sequences in the stain, whereas the other chromosomes would only be stained by the shared sequences. Thus, only a modest decrease in hybridization of the shared sequences in the stain would substantially enhance the contrast. Thus contaminating sequences which only hybridize to non-targeted sequences, for example, impurities in a library, can be tolerated in the stain to the extent that the sequences do not reduce the staining contrast below useful levels.

Detecting the 20q13 Amplicon.

As explained above, detection of amplification in the 20q13 amplicon is indicative of the presence and/or prognosis of a large number of cancers. These include, but are not limited to breast, ovary, bladder, head and neck, and colon.

In a preferred embodiment, a 20q13 amplification is detected through the hybridization of a probe of this invention to a target nucleic acid (*e.g.* a chromosomal sample) in which it is desired to screen for the amplification. Suitable hybridization formats are well known to those of skill in the art and include, but are not limited to,

BEST AVAILABLE COPY

variations of Southern Blots, *in situ* hybridization and quantitative amplification methods such as quantitative PCR (*see, e.g. Sambrook, supra., Kallioniemi et al., Proc. Natl Acad Sci USA*, 89: 5321-5325 (1992), and *PCR Protocols, A Guide to Methods and Applications*, Innis *et al.*, Academic Press, Inc. N.Y., (1990)).

In situ Hybridization.

In a preferred embodiment, the 20q13 amplicon is identified using *in situ* hybridization. Generally, *in situ* hybridization comprises the following major steps: (1) fixation of tissue or biological structure to be analyzed; (2) prehybridization treatment of the biological structure to increase accessibility of target DNA, and to reduce nonspecific binding; (3) hybridization of the mixture of nucleic acids to the nucleic acid in the biological structure or tissue; (4) posthybridization washes to remove nucleic acid fragments not bound in the hybridization and (5) detection of the hybridized nucleic acid fragments. The reagent used in each of these steps and their conditions for use vary depending on the particular application.

In some applications it is necessary to block the hybridization capacity of repetitive sequences. In this case, human genomic DNA is used as an agent to block such hybridization. The preferred size range is from about 200 bp to about 1000 bases, more preferably between about 400 to about 800 bp for double stranded, nick translated nucleic acids.

Hybridization protocols for the particular applications disclosed here are described in Pinkel *et al. Proc. Natl. Acad. Sci. USA*, 85: 9138-9142 (1988) and in EPO Pub. No. 430,402. Suitable hybridization protocols can also be found in *Methods in Molecular Biology Vol. 33: In Situ Hybridization Protocols*, K.H.A. Choo, ed., Humana Press, Totowa, New Jersey, (1994). In a particularly preferred embodiment, the hybridization protocol of Kallioniemi *et al., Proc. Natl Acad Sci USA*, 89: 5321-5325 (1992) is used.

Typically, it is desirable to use dual color FISH, in which two probes are utilized, each labelled by a different fluorescent dye. A test probe that hybridizes to the region of interest is labelled with one dye, and a control probe that hybridizes to a different region is labelled with a second dye. A nucleic acid that hybridizes to a stable portion of the chromosome of interest, such as the centromere region, is often most

BEST AVAILABLE COPY

useful as the control probe. In this way, differences between efficiency of hybridization from sample to sample can be accounted for.

The FISH methods for detecting chromosomal abnormalities can be performed on nanogram quantities of the subject nucleic acids. Paraffin embedded tumor sections can be used, as can fresh or frozen material. Because FISH can be applied to the limited material, touch preparations prepared from uncultured primary tumors can also be used (*see, e.g., Kallioniemi, A. et al., Cytogenet. Cell Genet.* 60: 190-193 (1992)). For instance, small biopsy tissue samples from tumors can be used for touch preparations (*see, e.g., Kallioniemi, A. et al., Cytogenet. Cell Genet.* 60: 190-193 (1992)). Small numbers of cells obtained from aspiration biopsy or cells in bodily fluids (*e.g., blood, urine, sputum and the like*) can also be analyzed. For prenatal diagnosis, appropriate samples will include amniotic fluid and the like.

Southern Blots

In a Southern Blot, a genomic or cDNA (typically fragmented and separated on an electrophoretic gel) is hybridized to a probe specific for the target region. Comparison of the intensity of the hybridization signal from the probe for the target region (*e.g., 20q13*) with the signal from a probe directed to a control (non amplified) such as centromeric DNA, provides an estimate of the relative copy number of the target nucleic acid.

Kits Containing 20q13 Amplicon Probes.

This invention also provides diagnostic kits for the detection of chromosomal abnormalities at 20q13. In a preferred embodiment, the kits include one or more probes to the 20q13 amplicon described herein. The kits can additionally include blocking probes, instructional materials describing how to use the kit contents in detecting 20q13 amplicons. The kits may also include one or more of the following: various labels or labeling agents to facilitate the detection of the probes, reagents for the hybridization including buffers, a metaphase spread, bovine serum albumin (BSA) and other blocking agents, sampling devices including fine needles, swabs, aspirators and the like, positive and negative hybridization controls and so forth.

EXAMPLES

BEST AVAILABLE COPY

The following examples are offered to illustrate, but not to limit the present invention.

Example 1

PROGNOSTIC IMPLICATIONS OF AMPLIFICATION OF CHROMOSOMAL REGION 20q13 IN BREAST CANCER

Patients and tumor material.

Tumor samples were obtained from 152 women who underwent surgery for breast cancer between 1987 and 1992 at the Tampere University or City Hospitals. One hundred and forty-two samples were from primary breast carcinomas and 11 from metastatic tumors. Specimens from both the primary tumor and a local metastasis were available from one patient. Ten of the primary tumors that were either *in situ* or mucinous carcinomas were excluded from the material, since the specimens were considered inadequate for FISH studies. Of the remaining 132 primary tumors, 128 were invasion ductal and 4 lobular carcinomas. The age of the patients ranged from 29 to 92 years (mean 61). Clinical follow-up was available from 129 patients. Median follow-up period was 45 months (range 1.4-177 months). Radiation therapy was given to 77 of the 129 patients (51 patients with positive and 26 with negative lymph nodes), and systemic adjuvant therapy to 36 patients (33 with endocrine and 3 with cytotoxic chemotherapy). Primary tumor size and axillary node involvement were determined according to the tumor-node metastasis (TNM) classification. The histopathological diagnosis was evaluated according to the World Health Organization (11). The carcinomas were graded on the basis of the tubular arrangement of cancer cells, nuclear atypia, and frequency of mitotic or hyperchromatic nuclear figures according to Bloom and Richardson, *Br. J. Cancer*, 11: 359-377 (1957).

Surgical biopsy specimens were frozen at -70°C within 15 minutes of removal. Cryostat sections (5-6 μ m) were prepared for intraoperative histopathological diagnosis, and additional thin sections were cut for immunohistochemical studies. One adjacent 200 μ m thick section was cut for DNA flow cytometric and FISH studies.

Cell preparation for FISH.

After histological verification that the biopsy specimens contained a high proportion of tumor cells, nuclei were isolated from 200 μ m frozen sections according to a modified Vindelov procedure for DNA flow cytometry, fixed and dropped on slides for

BEST AVAILABLE COPY

FISH analysis as described by Hyytinen *et al.*, *Cytometry* 16: 93-99 (1994). Foreskin fibroblasts were used as negative controls in amplification studies and were prepared by harvesting cells at confluency to obtain G1 phase enriched interphase nuclei. All samples were fixed in methanol-acetic-acid (3:1).

Probes.

Five probes mapping to the 20q13 region were used (*see Stokke, et al., Genomics*, 26: 134-137 (1995)). The probes included Pl-clones for melanocortin-3-receptor (probe MC3R, fractional length from p-arm telomere (Flpter 0.81) and phosphoenolpyruvate carboxy kinase (PCK, Flpter 0.84), as well as anonymous cosmid clones RMC20C026 (Flpter 0.79). In addition, RMC20C001 (Flpter 0.825) and RMC20C030 (Flpter 0.85) were used. Probe RMC20C001 was previously shown to define the region of maximum amplification (Tanner *et al.*, *Cancer Res*, 54: 4257-4260 (1994)). One cosmid probe mapping to the proximal p-arm, RMC20C038 (FLpter 0.237) was used as a chromosome-specific reference probe. Test probes were labeled with biotin-14-dATP and the reference probe with digoxigenin-11-dUTP using nick translation (Kallioniemi *et al.*, *Proc. Natl Acad Sci USA*, 89: 5321-5325 (1992)).

Fluorescence in situ hybridization.

Two-color FISH was performed using biotin-labeled 20q13-specific probes and digoxigenin-labelled 20p reference probe essentially as described (*Id.*). Tumor samples were postfixed in 4% paraformaldehyde/phosphate-buffered saline for 5 min at 4 C prior to hybridization, dehydrated in 70%, 85% and 100% ethanol, air dried, and incubated for 30 min at 80°C. Slides were denatured in a 70% formamide/2x standard saline citrate solution at 72-74°C for 3 min, followed by a proteinase K digestion (0.5 µg/ml). The hybridization mixture contained 18 ng of each of the labeled probes and 10 µg human placental DNA. After hybridization, the probes were detected immunochemically with avidin-FITC and anti-digoxigenin Rhodamine. Slides were counterstained with 0.2 µM 4,6-diamidino-2-phenylindole (DAPI) in an antifade solution.

Fluorescence microscopy and scoring of signals in interphase nuclei.

A Nikon fluorescence microscope equipped with double band-pass filters (Chromatechnology, Brattleboro, Vermont, USA) and 63 x objective (NA 1.3) was used

for simultaneous visualization of FITC and Rhodamine signals. At least 50 non-overlapping nuclei with intact morphology based on the DAPI counterstaining were scored to determine the number of test and reference probe hybridization signals.

Leukocytes infiltrating the tumor were excluded from analysis. Control hybridizations to normal fibroblast interphase nuclei were done to ascertain that the probes recognized a single copy target and that the hybridization efficiencies of the test and reference probes were similar.

The scoring results were expressed both as the mean number of hybridization signals per cell and as mean level of amplification (= mean of number of signals relative to the number of reference probe signals).

DNA flow cytometry and steroid receptor analyses.

DNA flow cytometry was performed from frozen 200 μ m sections as described by Kallioniemi, *Cytometry* 9: 164-169 (1988). Analysis was carried out using an EPICS C flow cytometer (Coulter Electronics Inc., Hialeah, Florida, USA) and the MultiCycle program (Phoenix Flow Systems, San Diego, California, USA). DNA-index over 1.07 (in over 20% of cells) was used as a criterion for DNA aneuploidy. In DNA aneuploid histograms, the S-phase was analyzed only from the aneuploid clone. Cell cycle evaluation was successful in 86% (108/126) of the tumors.

Estrogen (ER) and progesterone (PR) receptors were detected immunohistochemically from cryostat sections as previously described (17). The staining results were semiquantitatively evaluated and a histoscore greater than or equal to 100 was considered positive for both ER and PR (17).

Statistical Methods.

Contingency tables were analyzed with Chi square test for trend. Association between S-phase fraction (continuous variable) and 20q13 amplification was analyzed with Kruskal-Wallis test. Analysis of disease-free survival was performed using the BMDPIL program and Mautel-Cox test and Cox's proportional hazards model (BMDP2L program) was used in multivariate regression analysis (Dixon *BMDP Statistical Software*. London, Berkeley, Los Angeles: University of California Press, (1981)).

BEST AVAILABLE COPY

Amplification of 20q13 in primary breast carcinomas by fluorescence *in situ* hybridization.

The minimal region probe RMC20C001 was used in FISH analysis to assess the 20q13 amplification. FISH was used to analyze both the total number of signals in individual tumor cells and to determine the mean level of amplification (mean copy number with the RMC20C001 probe relative to a 20p-reference probe). In addition, the distribution of the number of signals in the tumor nuclei was also assessed. Tumors were classified into three categories: no, low and high level of amplification. Tumors classified as not amplified showed less than 1.5 fold-copy number of the RMC20C001 as compared to the p-arm control. Those classified as having low-level amplification had 1.5-3-fold average level of amplification. Tumors showing over 3-fold average level of amplification were classified as highly amplified.

The highly amplified tumors often showed extensive intratumor heterogeneity with up to 40 signals in individual tumor cells. In highly amplified tumors, the RMC20C001 probe signals were always arranged in clusters by FISH, which indicates location of the amplified DNA sequences in close proximity to one another *e.g.* in a tandem array. Low level 20q13 amplification was found in 29 of the 132 primary tumors (22%), whereas nine cases (6.8%) showed high level amplification. The overall prevalence of increased copy number in 20q13 was thus 29% (38/132).

Defining the minimal region of amplification.

The average copy number of four probes flanking RMC20C001 was determined in the nine highly amplified tumors. The flanking probes tested were melanocortin-3-receptor (MC3R, FLpter 0.81), phosphoenolpyruvate carboxykinase (PCK, 0.84), RMC20C026 (0.79) and RMC20C030 (0.85). The amplicon size and location varied slightly from one tumor to another but RMC20C001 was the only probe consistently highly amplified in all nine cases.

Association of 20q13 amplification with pathological and biological features.

The 20q13 amplification was significantly associated with high histologic grade of the tumors ($p=0.01$). This correlation was seen both in moderately and highly amplified tumors (Table 1). Amplification of 20q13 was also significantly associated

with aneuploidy as determined by DNA flow cytometry ($p=0.01$, Table 1.) The mean cell proliferation activity, measured as the percentage of cells in the S-phase fraction, increased ($p=0.0085$ by Kruskal-Wallis test) with the level of amplification in tumors with no, low and high levels of amplification (Table 1). No association was found with the age of the patient, primary tumor size, axillary nodal or steroid hormone-receptor status (Table 1).

Table 1. Clinicopathological correlations of amplification at chromosomal region 20q13 in 132 primary breast cancers.

Pathobiologic feature	20q13 amplification status			p-value ¹
	NO Number of patients (%)	LOW LEVEL Number of patients (%)	HIGH LEVEL Number of patients (%)	
All primary tumors	94 (71%)	29 (22%)	9 (6.8%)	
Age of patients				
< 50 years	17 (65%)	6 (23%)	3 (12%)	.39
≥ 50 years	77 (73%)	23 (22%)	6 (5.7%)	
Tumor size				
< 2 cm	33 (79%)	7 (17%)	2 (4.8%)	.16
≥ 2 cm	58 (67%)	22 (25%)	7 (8.0%)	
Nodal status				
Negative	49 (67%)	19 (26%)	5 (6.8%)	.41
Positive	41 (75%)	10 (18%)	4 (7.3%)	
Histologic grade				
I - II	72 (76%)	18 (19%)	5 (5.3%)	.01
III	16 (52%)	11 (35%)	4 (13%)	
Estrogen receptor status				
Negative	30 (67%)	10 (22%)	5 (11%)	.42
Positive	59 (72%)	19 (23%)	4 (4.9%)	
Progesterone receptor status				
Negative	57 (69%)	20 (24%)	6 (7.2%)	.53
Positive	32 (74%)	8 (19%)	3 (7.0%)	
DNA ploidy				
Diploid	45 (82%)	8 (14.5%)	2 (3.6%)	.01
Aneuploid	44 (62%)	20 (28%)	7 (10%)	
S-phase fraction (%)	mean ± SD 9.9 ± 7.2	mean ± SD 12.6 ± 6.7	mean ± SD 19.0 ± 10.5	.0085 ¹

Kruskal-Wallis Test.

Relationship between 20q13 amplification and disease-free survival.

Disease-free survival of patients with high-level 20q13 amplification was significantly shorter than for patients with no or only low-level amplification ($p=0.04$).

BEST AVAILABLE COPY

Disease-free survival of patients with moderately amplified tumors did not differ significantly from that of patients with no amplification. Among the node-negative patients (n=79), high level 20q13 amplification was a highly significant prognostic factor for shorter disease-free survival ($p=0.002$), even in multivariate Cox's regression analysis ($p=0.026$) after adjustment for tumor size ER, PR grade, ploidy and S-phase fraction.

20q13 amplification in metastatic breast tumors.

Two of 11 metastatic breast tumors had low level and one high level 20q13 amplification. Thus, the overall prevalence (27%) of increased 20q13 copy number in metastatic tumors was a similar to that observed in the primary tumors. Both a primary and a metastatic tumor specimens were available from one of the patients. This 29-year old patient developed a pectoral muscle infiltrating metastasis eight months after total mastectomy. The patient did not receive adjuvant or radiation therapy after mastectomy. The majority of tumor cells in the primary tumor showed a low level amplification, although individual tumor cells (less than 5% of total) contained 8-20 copies per cell by FISH. In contrast, all tumor cells from metastasis showed high level 20q13 amplification (12-50 copies per cell). The absolute copy number of the reference probe remained the same suggesting that high level amplification was not a result of an increased degree of aneuploidy.

Diagnostic and Prognostic Value of the 20q13 Amplification.

The present findings suggest that the newly-discovered 20q13 amplification may be an important component of the genetic progression pathway of certain breast carcinomas. Specifically, the foregoing experiments establish that: 1) High-level 20q13 amplification, detected in 7% of the tumors, was significantly associated with decreased disease-free survival in node-negative breast cancer patients, as well as with indirect indicators of high-malignant potential, such as high grade and S-phase fraction. 2) Low-level amplification, which was much more common, was also associated with clinicopathological features of aggressive tumors, but was not prognostically significant. 3) The level of amplification of RMC20C001 remains higher than amplification of nearby candidate genes and loci indicating that a novel oncogene is located in the vicinity of RMC20C001.

High-level 20q13 amplification was defined by the presence of more than 3-fold higher copy number of the 20q13 amplification is somewhat lower than the amplification frequencies reported for some of the other breast cancer oncogenes, such as ERBB2 (17q12) and Cyclin-D (11q13) (Borg *et al.*, *Oncogene*, 6: 137-143 (1991), Van de Vijver *et al.* *Adv. Canc. Res.*, 61: 25-56 (1993)). However, similar to what has been previously found with these other oncogenes (Swab, *et al.*, *Genes Chrom. Canc.*, 1: 181-193 (1990), Borg *et al.*, *supra.*), high-level 20q13 amplification was more common in tumors with high grade or high S-phase fraction and in cases with poor prognosis. Although only a small number of node-negative patients was analyzed, our results suggest that 20q13 amplification might have independent role as a prognostic indicator. Studies to address this question in large patient materials are warranted. Moreover, based on these survival correlations, the currently unknown, putative oncogene amplified in this locus may confer an aggressive phenotype. Thus, cloning of this gene is an important goal. Based on the association of amplification with highly proliferative tumors one could hypothesize a role for this gene in the growth regulation of the cell.

The role of the low-level 20q13 amplification as a significant event in tumor progression appears less clear. Low-level amplification was defined as 1.5-3-fold increased average copy number of the 20q13 probe relative to the p-arm control. In addition, these tumors characteristically lacked individual tumor cells with very high copy numbers, and showed a scattered, not clustered, appearance of the signals. Accurate distinction between high and low level 20q13 amplification can only be reliably done by FISH, whereas Southern and slot blot analyses are likely to be able to detect only high-level amplification, in which substantial elevation of the average gene copy number takes place. This distinction is important, because only the high amplified tumors were associated with adverse clinical outcome. Tumors with low-level 20q13 amplification appeared to have many clinicopathological features that were in between of those found for tumors with no and those with high level amplification. For example, the average tumor S-phase fraction was lowest in the non-amplified tumors and highest in the highly amplified tumors. One possibility is that low-level amplification precedes the development of high level amplification. This has been shown to be the case, *e.g.*, in the development of drug resistance-gene amplification *in vitro* (Stark, *Adv. Canc. Res.*, 61: 87-113 (1993)). Evidence supporting this hypothesis was found in one of our

BEST AVAILABLE COPY

patients, whose local metastasis contained a much higher level of 20q13 amplification than the primary tumor operated 8 months earlier.

Finally, our previous paper reported a 1.5 Mb critical region defined by RMC20C001 probe and exclusion of candidate genes in breast cancer cell lines and in a limited number of primary breast tumors. Results of the present study confirm these findings by showing conclusively in a larger set of primary tumors that the critical region of amplification is indeed defined by this probe.

The present data thus suggest that the high-level 20q13 amplification may be a significant step in the progression of certain breast tumors to a more malignant phenotype. The clinical and prognostic implications of 20q13 amplification are striking and location of the minimal region of amplification at 20q13 has now been defined.

It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference for all purposes.

BEST AVAILABLE COPY

WHAT IS CLAIMED IS:

1 1. A method of detecting a chromosome abnormality at about
2 position FLpter 0.825 on human chromosome 20, the method comprising:
3 contacting a chromosome sample from a patient with a
4 composition consisting essentially of one or more labeled nucleic acid probes
5 each of which binds selectively to a target polynucleotide sequence at about
6 position FLpter 0.825 on human chromosome 20 under conditions in which the
7 probe forms a stable hybridization complex with the target sequence; and
8 detecting the hybridization complex.

1 2. The method of claim 1, wherein the probe comprises a
2 nucleic acid that specifically hybridizes under stringent conditions to a nucleic
3 acid selected from a group consisting of the nucleic acids listed in Table 1, and
4 Table 2.

1 3. The method of claim 2, wherein the probe is selected from
2 a group consisting of the probes listed in Table 1 and Table 2.

1 4. The method of claim 1, wherein the step of detecting the
2 hybridization complex comprises determining the copy number of the target
3 sequence.

1 5. The method of claim 1, wherein the probe is labeled with
2 digoxigenin or biotin.

1 6. The method of claim 1, wherein the hybridization complex
2 is detected in interphase nuclei in the sample.

1 7. The method of claim 1, wherein the chromosome
2 abnormality is an amplification.

1 8. The method of claim 1, wherein the probes are labeled
2 with digoxigenin or biotin.

BEST AVAILABLE COPY

1 9. The method of claim 1, wherein the step of detecting the
2 hybridization complexes is carried out by detecting a fluorescent label.

1 10. The method of claim 9, wherein the fluorescent label is
2 FITC.

1 11. The method of claim 1, further comprising contacting the
2 sample with a reference probe which binds selectively to a chromosome 20
3 centromere.

1 12. A composition comprising a labeled nucleic acid probe
2 which binds selectively to a target polynucleotide sequence at about FLpter
3 0.825 on human chromosome 20.

1 13. The composition of claim 12, wherein the probe is
2 labelled with digoxigenin or biotin.

1 14. The composition of claim 1, wherein the probe comprises
2 a nucleic acid that hybridizes under stringent conditions to one or more nucleic
3 acids selected from the group consisting of the nucleic acids listed in Table 1
4 and Table 2.

1 15. The composition of claim 14, wherein the probe comprises
2 one or more nucleic acids selected from the group consisting of the nucleic
3 acids listed in Table 1 and Table 2.

1 16. A kit for the detection of a chromosomal abnormality at
2 about position FLpter 0.825 on human chromosome 20, the kit comprising a
3 compartment which contains a labeled nucleic acid probe which binds
4 selectively to a target polynucleotide sequence at about FLpter 0.825 on human
5 chromosome 20.

1 17. The kit of claim 16, wherein label is selected from the
2 group consisting of digoxigenin and biotin.

1 18. The composition of claim 1, wherein the probe is one or
2 more probes selected from the group consisting of the nucleic acids of Table 1,
3 and Table 2.

1 19. The kit of claim 16, further comprising a reference probe
2 specific to a sequence in the centromere of chromosome 20.

1 20. The kit of claim 16, further comprising Texas red avidin
2 and biotin-labeled goat anti-avidin antibodies.

BEST AVAILABLE COPY

1/4




FIG. 1A.




FIG. 1B.

SUBSTITUTE SHEET (RULE 26)

2/4





FIG. 2.

BEST AVAILABLE COPY

SUBSTITUTE SHEET (RULE 26)



BEST AVAILABLE COPY




FIG. 4.

INTERNATIONAL SEARCH REPORT

Inter. application No.
PCT/US96/16085

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12Q 1/68; C07H 21/04

US CL : 435/6; 536/24.3

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6; 536/24.3

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN searched, uspat, medline, biosis, caplus

search terms: detect chromosome abnormality, FLpter 0.825, fluorescent labeled, hubridization complex

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y, P	US 5,472,842 A (STOKKE et al.) 05 December 1995, column 1, lines 66-67; column 2, lines 1-45.	1-20
X	TANNER et al. Increased Copy Number at 20q13 in Breast Cancer: Defining the Critical Region and Exclusion of Candidate Genes. Cancer Research. 15 August 1994, Vol. 54, pages 4257-4260, especially page 4257, abstract.	1, 4-13, 16, 17, 19, 20
Y	GYAPAY et al. The 1993-1994 genethon human genetic linkage map. Nature Genetics. June 1994, Vol. 7. pages 246-339, especially page 329.	2, 3, 14, 15, 18
Y	EP 0 571 911 A2 (BECTON, DICKINSON & COMPANY) 01 December 1993, page 14, claim 3.	2, 3, 14, 15, 18

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

26 DECEMBER 1996

Date of mailing of the international search report

30 JAN 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JOYCE TUNG

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)*

BEST AVAILABLE COPY

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/16085

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Datebase EST-STS on MPsrch Accession No. G08049, Murray et al. 'Cooperative human linkage center' 08 August 1995.	2, 3, 14, 15, 18

Form PCT/ISA/210 (continuation of second sheet)(July 1992)*

BEST AVAILABLE COPY